



ISSN (P): 2617-7226
ISSN (E): 2617-7234
www.patholjournal.com
2020; 3(2): 37-43
Received: 22-02-2020
Accepted: 24-03-2020

Nanigopal Bhattacharya
Professor & H.O.D,
Department of Pathology,
Vivekananda Institute of
Medical Sciences, Kolkata,
West Bengal, India

Shaheena Perween
Junior resident & P.G.T ,
Department of Pathology,
Vivekananda Institute of
Medical Sciences, Kolkata,
West Bengal, India

Krishnendu Gupta
Professor , Department of
Obs. & Gynaecology,
Vivekananda Institute of
Medical Sciences, Kolkata,
West Bengal, India

Corresponding Author:
Nanigopal Bhattacharya
Professor & H.O.D,
Department of Pathology,
Vivekananda Institute of
Medical Sciences, Kolkata,
West Bengal, India

Histopathological study of surgically resected specimens of fallopian tube

Nanigopal Bhattacharya, Shaheena Perween and Krishnendu Gupta

DOI: <https://doi.org/10.33545/pathol.2020.v3.i2a.222>

Abstract

Background: Fallopian Tubes are very common surgical specimen received in the pathological laboratory. It is affected by wide spectrum of both non neoplastic and neoplastic lesions. But there is very few studies described the frequency of various histopathological spectrums of lesions of fallopian tube.

Aims & Objectives: To study the various histopathological lesions seen in surgically resected specimens of fallopian tube and their age distributions.

Material and Method: A total of 200 surgically resected specimen of fallopian tube of all age group of patients which are received for one and half year duration, are processed with routine histopathological techniques. Histopathological features were studied on Haematoxylin and Eosin stain.

Results: In our studies majority of the cases are belongs to 40 to 60 years of age out of 200 specimens. In this study, most of the patients were undergone total abdominal hysterectomy with salpingo-oophorectomy (60%).

In our study out of 200 cases 63 (31.50%) specimens show unremarkable histology. Only 1(0.50%) case shows neoplastic pathology. Most common histopathological findings observed are paratubal cyst, ectopic tubal gestation and Walthard Cell Nests.

Conclusion: Present study shows broad spectrum of pathological lesions in fallopian tube. Majority of the lesions are non neoplastic in nature. Thorough histopathological examination of fallopian tube specimens is necessary to document many lesions including rare primary intraepithelial malignancy which can be missed clinically and found incidentally on histopathological examination.

Keywords: Fallopian tube, histopathology, non neoplastic lesion, neoplastic lesion

Introduction

The fallopian tubes are complex structures that connect ovaries to the endometrial cavity. They are sites of various interactions necessary for normal pregnancy ^[1]. It is a common specimen in a pathology laboratory and may be examined either alone as a salpingectomy specimen or as a part of a more complex specimen from a hysterectomy and/or oophorectomy operation ^[2].

Although fallopian tube is affected by a wide spectrum of diseases, literature search reveals that there are only few studies documenting histological changes in fallopian tube removed for all reasons. One of the causes of secondary infertility is inflammation and neoplasm of fallopian tube. Ectopic pregnancy even though uncommon, but may be detected by routine ultrasound, which if undetected could cause maternal death ^[3, 4]. Ectopic pregnancy is a complication of pregnancy in which the embryo is attached outside the uterus ^[8], and fallopian is a common site. Signs and symptoms classify include abdominal pain and vaginal bleeding ^[9]. Those who have previously had an ectopic pregnancy are at much higher risk of having another one ^[10]. Most of the ectopic pregnancies (90%) occur on the fallopian tube, which are known as tubal pregnancy ^[10]. The rate of ectopic pregnancy is about 1% to 2% that of live birth in developed countries, though it may be as high as 4% among those using assisted reproductive technology ^[8].

Inflammatory disease of tube (salpingitis) remains responsible for a significant percentage of cases of secondary infertility by occlusion or stenosis. Hence in this modern era of advancement it has become a necessity to study in details the pathology of various fallopian tube diseases. Risk factors for ectopic pregnancy also include pelvic inflammatory disease, often due to tuberculosis, salpingitis, prior tubal surgery and the uses of assisted reproductive technology ^[10].

Bacterial infection of the fallopian tube is a common disease, and its incidence keeps increasing. It may follow invasive procedures (such as curettage or the insertion of intrauterine devices^[11]), and it commonly accompanies endometriosis^[12], but in most cases it is due to an ascending infection, often sexually transmitted^[13]. The inflammation may result in fusion of tubal plicae and obliteration of the ostium. Obstruction of the fimbrial end of fallopian tube and, less commonly, of the intramural or isthmic portion may lead to infertility^[14]. Microscopically, the trapped epithelial spaces produce a complex gland like pattern that can stimulate malignancy^[15].

The lumen is often distended and filled with secretions or pus (pyosalpinx). Massive intraluminal hemorrhage may lead to the formation of hematosalpinx, a rare condition that needs to be distinguished from ruptured tubal pregnancy. In chronic inflammatory cases, the tubal wall is markedly fibrotic and serosal adhesion is prominent. The inflammatory exudates often spread to the ovary, resulting in the formation of a tubo-ovarian abscess and obliteration of the pelvic anatomic relationships. The rupture of such an abscess may lead to localized or generalized peritonitis and requires prompt surgical intervention^[16, 17]. Hydrosalpinx is generally regarded as the end stage of a purulent salpingitis in which pus has been reabsorbed and replaced by a transudate of plasma^[18]. Pelvic inflammatory diseases are the generic term used for the inflammatory process of this region in which the fallopian tube is the epicenter and presumably the source of inflammation^[19, 20, 21]. Eschenbach *et al.*^[22], recovered *Neisseria gonorrhoeae* from 91 of 204 cases of acute pelvic inflammatory disease in patients with cervical gonococcal inflammation. Chlamydial infection is also common, accounting for over 20% and perhaps as many as half of the cases^[23, 24]. This etiology should be suspected in cases of chronic salpingitis accompanied by marked lymphoid follicular hyperplasia^[23].

Tuberculosis of the fallopian tube develops by the haematogenous route. In advanced cases, both tubes are replaced by caseous tuberculous masses, approximately 80% of cases^[25]. Granulomatous inflammation of the tubes can also be produced by schistosoma, oxyuris vermicularis, actinomyces and other organism^[26]. Sarcoidosis and Crohn's disease may be accompanied by tubal involvement. Foreign bodies introduced for diagnostic or therapeutic measures can induce bizarre granulomatous responses. Reaction to the lipoidal that used to be introduced in the Rubin's test may be so proliferative as to resemble a neoplasm. The push of a uterine sound may drive lubricant into the tube, causing lipoid granulomas^[27].

Xanthogranulomatous salpingitis is characterized by infiltrate of foamy histiocytes. Some authors make distinction between Xanthogranulomatous and Pseudoxanthomatous salpingitis is probably unwarranted^[28]. Accordingly, some authors refer to it as pseudoxanthomatous salpingitis^[29].

Giant cell arteritis is occasionally found in the tube, ovaries and uterus of postmenopausal patients, either as an isolated finding (most frequently) or as a manifestation of a generalized immune mediated disease^[30].

Torsion of the fallopian tube and ovary is usually secondary to inflammation or tumor, but occasionally it develops in a previously normal organ, the appearance at surgery being that of a hemorrhagic infarct. This phenomenon can occur in

adults^[31] as well as in infants and children^[32].

Walthard cell nests are small, glistening, round collection of flat to cuboidal cells with the appearance of urothelium located on the tubal serosa^[33], sometimes accompanied by cystic changes. They are probably of mesothelial rather than mullerian or wolffian nature. They should not be mistaken for serosal implants in patients with ovarian neoplasm.

Para tubal cysts, traditionally known as hydatid cyst of Morgagni, are commonly seen as small round cysts attached by a pedicle to the fimbrial end of the tube. Their wall is paper thin and their content is clear. Occasionally, they attain a large size and may undergo torsion^[34]. Most are lined by tubal columnar epithelium containing both ciliated and secretory cells, sometimes projecting in a papillary fashion into the lumen and covered by a thin layer of smooth muscle^[35]. Their appearance is consistent with origin from mullerian-type structure. Other paratubal cysts, lined by flat cells and surrounded by a thin fibrous wall, are regarded as of mesothelial origin^[36].

Endometriosis frequently involves the tube in the form of nodules located in the wall or serosa^[37]. Decidual reaction of the tubal mucosa is a common finding in specimens of tubal ligation obtained at the time of caesarean section; it appears as small nodular collections of decidual cells covered by a flattened, sometimes inflamed epithelium. Similar changes have been documented following hormonal therapy^[38].

Arias-stella reaction can occur in the fallopian tube epithelium in association with either orthotopic or tubal pregnancy^[39].

The fallopian tube primary malignancy is extremely rare and accounts for less than 0.14% to 0.18% of all genital malignancies with an annual incidence of 3.6 per million woman. Metastatic carcinoma of fallopian tube appears to be more common. The fallopian tube malignancy possesses a diagnostic challenge as it is difficult to distinguish between primary ovarian and primary endometrial carcinoma with primary fallopian tube carcinoma^[5, 6]. Serous tubal intraepithelial carcinoma (STIC) is rare pathologic finding at the time of benign gynecologic surgery. It arises in the distal fimbrial end of the fallopian tube and likely represents a precursor lesion to high grade pelvic serous carcinoma. A putative non malignant precursor to STIC, termed the p53 signature, has recently been described in fallopian tube^[7]. Traditionally primary carcinoma of the fallopian tube has been regarded as very rare, accounting for approximately 1% of primary genital tract malignancies^[40]. However, several recent studies following a standardized grossing protocol observed that it reaches approximately 15% of all adnexal tumors,^[41] including a fair number of early stages. The interesting possibility is that a high percentage of ovarian and primary peritoneal serous carcinomas may originate in the tubal fimbriae. Microscopically, all the major types of carcinoma known to occur in the ovary have been reported in the tube. The most common type is papillary serous carcinoma^[42]. Borderline epithelial neoplasm has also been described in the fallopian tube.

Fallopian tube is affected by a wide spectrum of diseases varying from salpingitis to carcinoma. As this is a common specimen in the surgical pathology laboratory, knowledge of the frequency of these various pathological lesions involving fallopian tube can be valuable to pathologists, who may

identify an unusual histological finding.

Material and Method

A total of 200 surgically resected specimen of fallopian tube of all age group of patients which are received for one and half year duration, are processed with routine histopathological techniques. All poorly fixed specimens are excluded from the study. Minimum of two cross sections from proximal, mid portions and one longitudinal section from distal fimbriated end of each fallopian tube are taken for examination. Histopathological features were studied on Haematoxylin and Eosin stain and all microscopic features are noted down.

Result

In our study the youngest patient was 11-20 years and eldest one was 71-80 years. Most of the patient's age is between 41-60 years. (Table.1) Maximum numbers of patient are from pre and post menopausal age group. However all cases of tubal gestation, maximum number of endometriosis and stromal decidual changes are from reproductive age group. Only two cases of granulomatous salpingitis are seen who are also from reproductive age (Table no.2). In our study most of the patients are in the post-menopausal age group (44%) and premenopausal age group (25.50%), Most of the patients undergone total abdominal hysterectomy with salpingo-oophorectomy (60%) and few patients undergone unilateral salpingectomy (10%). (Table.3)

In our study most of the cases show unremarkable histology i.e., 31.50% and least number of cases show serous tubal intraepithelial carcinoma, high grade (0.50%) (Table no.4) Most common histopathological findings observed are Para tubal cyst, ectopic tubal gestation and Walthard Cell Nests.

Discussion

The study conducted by Bagwan IN *et al.* (2004) [4] observed that majority of the cases belonged to the age group of 36 to 45 years. Gon S *et al.* (2013) [5] showed the same findings. In the present study majority cases are in the age group of 40 to 60 years.

According to the study conducted by Jennifer L. Hunt *et al.* (2002) [2] specimens of fallopian tubes received following tubal ligation and total abdominal hysterectomy with bilateral salpingo-oophorectomy in 38.62% of the cases each followed by salpingectomy and unilateral salpingo-oophorectomy in 22.76% of the cases. According to Bagwan IN *et al.* (2004) [4], most common surgical procedure performed was total abdominal hysterectomy with bilateral salpingo-oophorectomy in 72.77% of the cases followed by salpingo-oophorectomy in 22.9% cases and tubal ligation in 4.24% cases. According to Gon S *et al.* (2013) [5], total abdominal hysterectomy with bilateral salpingo-oophorectomy was done in 75.11% cases followed by salpingectomy in 14% cases, unilateral salpingo-oophorectomy in 5.86% cases and tubal ligation in 5% cases. Prachi k *et al.* [43] showed tubal ligation was done in 51.5% cases followed by total abdominal hysterectomy with bilateral salpingo-oophorectomy in 33% cases, salpingectomy in 9% cases, unilateral salpingo-oophorectomy in 3% cases, total abdominal hysterectomy with unilateral salpingo-oophorectomy in 2% cases and bilateral salpingo-oophorectomy in 1.5% cases.

In the present study, total abdominal hysterectomy with salpingo-oophorectomy done in 60% cases followed by unilateral salpingo-oophorectomy in 15% cases, unilateral salpingectomy in 10% and tubal ligation in 15% cases.

Most studies showed unremarkable histology. The present study also shows unremarkable pathology in a maximum number of cases (36.1%).

Study conducted by Jennifer L. Hunt *et al.* [2], observed that most common lesions were non-neoplastic, in which inflammatory lesions such as salpingitis was most common and seen in 32.5% of the specimens followed by tubal epithelial hyperplasia in 10.8% of the specimens. Bagwan IN *et al.* [4] also observed the most common lesions were non-neoplastic in which inflammatory pathology were the most common, seen in 18.34% cases. The inflammatory lesions were salpingitis (10.19%), hydrosalpinx (7.86%) and pyosalpinx (0.29%) of the fallopian tube.

Present study shows, chronic salpingitis 5% and chronic salpingitis with mucinous metaplasia in 1% of cases. Incidence of hydrosalpinx is 3% and hematosalpinx is 2% in this study.

Prachi *et al.* [43] and Bagwan IN *et al.* [4] found tubal epithelial hyperplasia in 1.34% and 0.29% of cases respectively. In our study, tubal epithelial hyperplasia is seen in 3% cases.

The study done by N Dahiya *et al.* [44], tubal ectopic pregnancy was commonly seen in the younger age group of 25-30 years. Shraddha Shetty K *et al.* [45], showed in 25-30 years and according to C. Li *et al.* [46], most common age group was 20-29 years. Our study shows 7.50% of tubal gestation in the age group of 21-30 years, which is almost same with other studies by N Dahiya *et al.* [44] and Shraddha Shetty K *et al.* [45].

Study done by Pratima Kujur *et al.* (2016) [47] showed 0.54% of tuberculous salpingitis cases. Lakshmi *et al.* [6] observed almost equal incidence of tuberculous salpingitis (0.59%).

Primary genital tuberculosis is quite rare. The most common site of genital tuberculosis is fallopian tubes, accounting to approximately 5%. In our study, granulomatous salpingitis is seen in 2 (1%) cases. Agarwal and Gupta⁴⁸ on their study of female genital tract found incidence of tuberculosis declining from 1.8% in 1974 to 0.8% in 1989. Epithelial tufting / Stratification / Hyperplasia (3.5%) are seen more with increased patient age. In our study it is in 3%. of specimens. Our study is almost same as Hunt JL, *et al.* (2002) [2]. Hunt JL, *et al.* (2002) [2]. also found lymphoid follicles in 2.1%. of cases. In our study it is 1%.

Pratima Kujur *et al.* (2016) [47] observed in their study, the incidence of paratubal cyst in 11.35% of patients. Our study shows incidence of paratubal cyst is 7%. In this study endometriosis is seen 5% of the cases. The Incidence of endometriosis is slightly higher as compared to other studies. (Pratima kujur, *et al.* 2016) [47]

Other non neoplastic lesions are seen in this study are decidual reaction (4%), Walthard Nests (10%) and Walthard Nests with cystic change (5%) which are slightly higher than other studies done by Prachi Kukreja *et al.* (2017) [43].

Incidence of paratubal cyst (Para mesonephric cyst 7% Mesonephric cyst 4% and Mesothelial cyst 5%) which is almost same as studies done by Prachi Kukreja *et al.* (2017) [43], Mehnaz Choudhury *et al.* (2019) [49] and Pratima Kujur *et al.* (2016) [47].

In all the studies, neoplastic lesions of fallopian tubes were rare finding. In our study also, neoplastic lesion such as serous tubal intraepithelial carcinoma (high grade) is found in only 0.5% of patient.

Financial Support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Table 1: shows distributions of total 200 cases according to their age-wise distribution of tubal lesions.

	11 - 20	21- 30	31- 40	41- 50	51- 60	61- 70	71 - 80	Total	Percentage
Chronic Salpingitis	0	0	1	4	3	1	1	10	5%
Chronic Salpingitis with Walthard Cell Nest	0	0	0	1	1	0	0	2	1%
Cystic Walthard Cell Nest	0	0	2	3	2	2	1	10	5%
Stromal Oedema	0	0	2	2	1	0	0	5	2.50%
Epithelial Hyperplasia	0	1	2	3	0	0	0	6	3%
Stromal Fibrosis	0	0	0	1	1	1	1	4	2%
Hydrosalpinx	0	1	1	3	1	0	0	6	3%
Granulomatous Salpingitis	0	1	1	0	0	0	0	2	1%
Tubal Gestation	2	9	4	0	0	0	0	15	7.50%
Stromal Decidual Change	2	4	1	1	0	0	0	8	4%
Lymphatic Dilatation and Presence of lymphoid Follicles	0	0	0	1	1	0	0	2	1%
Hematosalpinx	0	0	0	2	1	1	0	4	2%
Paramesonephric Cyst	0	1	1	6	3	3	0	14	7%
Mesonephric Cyst	0	1	2	3	1	1	0	8	4%
Mesothelial Cyst	0	1	2	1	5	1	0	10	5%
Endometriosis	1	2	2	2	2	1	0	10	5%
Serous Tubal Intraepithelial carcinoma- High grade	0	1	0	0	0	0	0	1	0.50%
Unremarkable	0	2	7	10	25	6	13	63	31.50%
Total								200	100%

Table 2: shows distribution of total 200 cases according to their menstrual status

DISTRIBUTION OF TUBAL LESIONS ON THE BASIS OF MENSTRUAL STATUS					
Fallopian Tube Pathology	Reproductive Age	Pre - Menopausal	Post - Menopausal	Total	% of Incidence
	11 - 40 yrs	40 - 50 yrs	50 - 80 yrs		
Chronic Salpingitis	1	4	5	10	5%
Chronic Salpingitis with Mucinous Metaplasia	0	1	1	2	1%
Walthard Cell Nest	4	8	8	20	10%
Cystic Walthard Cell Nest	2	3	5	10	5%
Stromal Oedema	2	2	1	5	2.50%
Epithelial Hyperplasia	3	3	0	6	3%
Stromal Fibrosis	0	1	3	4	2%
Hydrosalpinx	2	3	1	6	3%
Granulomatous Salpingitis	2	0	0	2	1%
Tubal Gestation	15	0	0	15	7.50%
Stromal Decidual Change	7	1	0	8	4%
Lymphatic Dilatation and Presence of Lymphoid Follicles	0	1	1	2	1%
Hematosalpinx	0	2	2	4	2%
Paramesonephric Cyst	2	6	6	14	7%
Mesonephric Cyst	3	3	2	8	4%
Mesothelial Cyst	3	1	6	10	5%
Endometriosis	5	2	3	10	5%
Serous Tubal Intraepithelial Carcinoma - High Grade	1	0	0	1	0.50%
Unremarkable	9	10	44	63	31.50%
TOTAL				200	100%

Table 3: shows distribution of total 200 cases according to types of surgery.

Type of Surgery	Total Number	Percentage
Total Abdominal Hysterectomy with Salpingo Oophorectomy	120	60%
Unilateral Salpingo - oophorectomy	30	15%
Unilateral Salpingectomy	20	10%
Tubal Ligation	30	15%
Total	200	100%

Table 4: shows distribution of total no of 200 cases according to their fallopian tube pathology.

Fallopian Tube Pathology	Total	% of Incidence
Chronic Salpingitis	10	5%
Chronic Salpingitis with Mucinous Metaplasia	2	1%
Walthard Cell Nest	20	10%
Cystic Walthard Cell Nest	10	5%
Stromal Oedema	5	2.50%
Epithelial Hyperplasia	6	3%
Stromal Fibrosis	4	2%
Hydrosalpinx	6	3%
Granulomatous Salpingitis	2	1%
Tubal Gestation	15	7.50%
Stromal Decidual Change	8	4%
Lymphatic Dilatation and Presence of Lymphoid Follicles	2	1%
Hematosalpinx	4	2%
Paramesonephric Cyst	14	7%
Mesonephric Cyst	8	4%
Mesothelial Cyst	10	5%
Endometriosis	10	5%
Serous Tubal Intraepithelial Carcinoma - High Grade	1	0.50%
Unremarkable	63	31.50%
Total	200	100%




Fig 1: Granulomatous Salpingitis H&E (40x)




Fig 4: Endometriosis H & E (10x)




Fig 2: Cystic Walthard Cell Nest H & E (20x)




Fig 5: Tubal Gestations H&E (20x)




Fig 3: Stromal Decidual Changes H & E (10x)




Fig 6: Mesothelial Cyst H&E (10x)




Fig 7: Paramesonephric Cyst H&E (20x)




Fig 8: Mesonephric CYST H & E (10x)




Fig 9: Serous Intraepithelial Carcinoma (Highgrade) H&E (20x)




Fig 10: Serous Intraepithelial Carcinoma (Highgrade) H&E (40x)

Conclusion

Present study shows broad spectrum of pathological lesions

in fallopian tube. Majority of the lesions are non neoplastic in nature. Neoplastic pathology is rare. Thorough histopathological examination of fallopian tube specimens is necessary to document many lesions including rare primary intraepithelial malignancy which can be missed clinically and found incidentally on histopathological examination.

Reference

1. Anderson MC. The fallopian tube. Symmers WS, editor. Systemic Pathology Female Reproductive System. London: Churchill Livingstone, 1991, 241-61.
2. Hunt JL, Lynn AA. Histologic features of surgically removed fallopian tubes. Arch Pathol Lab Med. 2002; 126(8):951-5
3. Patel J, Iyer RR. Spectrum of histopathological changes in fallopian tubes -a study of 350 cases. International journal of scientific research. 2016; 5(1):180-1.
4. Bhagwan IN, Harke AB, Malpani MR, Deshmukh MD. Histopathological study of spectrum of lesions encountered in the fallopian tube. J Obstet Gynecol Ind. 2004; 54(4):379-82.
5. Gon S, Basu A, Majumdar B, Das TK, Sengupta M, Ghosh D. Spectrum of histopathological lesions in the fallopian tubes. Journal of pathology of Nepal. 2013; 3:356-60.
6. Lakshmi K, Baleswari G, Mallikarjun C, Rao IB *et al.* Histopathological study of spectrum of lesions in the fallopian tubes. J Evol Med Dent Sci. 2015; 1:350-5.
7. Lee Y, Miron A, Drapkin R *et al.* A candidate precursor to serous carcinoma that originates in the distal fallopian tube. J Pathol. 2007; 211:26-35,
8. Kirk E, Bottomley C, Bourne T. "Diagnosing ectopic pregnancy and current concepts in the management of pregnancy of unknown location. Human Reproduction update. 2014; 20(2):250-61.
9. Crochet JR, Bastian LA, Chircau MV. "Dose this woman have an ectopic pregnancy ? the rational clinical examination systematic review". JAMA. 2013; 309(16):1722-9.
10. Cecchino GN, Araujo Junior E, Elito Junior J. Methotrexate for ectopic pregnancy: when and how". Archives of Gynecology and Obstetrics 2014; 290(3):417-23.
11. Charonis G, Larsson PG. Prolonged use of intrauterine contraceptive device as a risk factor for tubo-ovarian abscess. Acta Obstet Gynecol Scand 2009; 88:680-684
12. Seidman JD, Sherman ME, Bell KA, Katabuchi H, O'Leary TJ, Kurman RJ. Salpingitis, salpingoliths, and serous tumors of the ovaries: is there a connection? Int J Gynecol Pathol. 2002; 21:101-107.
13. Thor AD, Young RH, Clement PB. Pathology of the fallopian tube, broad ligament, peritoneum, and pelvic soft tissues. Hum Pathol. 1991; 22: 856-867.
14. Fortier KJ, Haney AF. The pathologic spectrum of uterotubal junction obstruction. Obstet Gynecol. 1985; 65:93-98
15. Cheung ANY, Young RH, Scully RE. Pseudocarcinomatous hyperplasia of the fallopian tube associated with salpingitis. A report of 14 cases. Am J Surg Pathol. 1994; 18:1125-1130
16. Mickal A, Sellmann AH, Beebe JL. Ruptured tubo ovarian abscess. Am J Obstet Gynecol 1968; 100:432-436.

17. Pedowitz R, Bloomfield RD. Ruptured adnexal abscess (tuboovarian) with generalized peritonitis. *Am J Obstet Gynecol.* 1964; 88:721-729.
18. David A, Garcia CS, Czernobilsky B. Human hydrosalpinx. Histologic study and chemical composition of fluid. *Am J Obstet Gynecol* 1969; 105:400-411
19. McCormack WM. Pelvic inflammatory disease. *N Engl J Med* 1994; 330:115-119.
20. Paavonen J. Pelvic inflammatory disease. From diagnosis to prevention. *Dermatol Clin* 1998; 16:747-756
21. Washington AE, Aral SO, Wolner-Hanssen P, Grimes DA, Holmes KK. Assessing risk for pelvic inflammatory disease and its sequelae. *JAMA* 1991; 266:2581-2586.
22. Eschenbach DA, Buchanan TM, Pollock HM, Forsyth PS, Alexander ER, Lin JS, Wang SP *et al.* Polymicrobial etiology of acute pelvic inflammatory disease. *N Engl J Med.* 1975; 29:166-17
23. Wallace TM, Hart WR. Acute chlamydial salpingitis with ascites and adnexal mass simulating a malignant neoplasm. *Int J Gynecol Pathol.* 1991; 10:394-401.
24. Winkler B, Reumann W, Mitao M, Gallo L, Richart RM, Crum CP. Immunoperoxidase localization of chlamydial antigens in acute salpingitis. *Am J Obstet Gynecol* 1985; 152:275-278
25. Nogales-Ortiz F, Tarancón I, Nogales FF Jr. The pathology of female genital tuberculosis. *Obstet Gynecol* 1979; 53:422-428
26. Erthan Y, Zekioglu O, Ozdemir N, Sen S. Unilateral salpingitis due to enterobius vermicularis. *Int J Gynecol Pathol.* 2000; 19:188-189.
27. Elliott GB, Brody H, Elliott KA. Implications of 'lipoid salpingitis'. *Fertil Steril.* 1965; 16:541-548.
28. Kostopoulou E, Daponte A, Kallitsaris A, Papamichali R, Kalodimos G, Messinis IE *et al.* Xanthogranulomatous salpingitis: report of three cases and comparison with a case of pseudoxanthomatous salpingitis. *ClinExp Obstet Gynecol* 2008; 35:291-294.
29. Seidman JD, Oberer S, Bitterman P, Aisner SC. Pathogenesis of pseudoxanthomatous salpingitis. *Mod Pathol.* 1993; 6:53-55.
30. Onuma K, Chu CT, Dabbs GJ. Asymptomatic giant-cell (temporal) arteritis involving the bilateral adnexa: case report and literature review. *Int J Gynecol Pathol.* 2007; 26:352-355.
31. Hansen OH. Isolated torsion of the fallopian tube. *Acta Obstet Gynecol Scand.* 1970; 49:3-6.
32. James DF, Barber HRK, Graber EA. Torsion of normal uterine adnexa in children. Report of three cases. *Obstet Gynecol.* 1970; 35:226-230.
33. Doss BJ, Jacques SM, Qureshi F, Ramirez NC, Lawrence WD. Extratubal secondary trophoblastic implants: clinicopathologic correlation and review of the literature. *Hum Pathol.* 1998; 29:184-187.
34. Goldrath MH, Platt LD. Treatment of ectopic tubal pregnancies by laparoscopy. *J Am Assoc Gynecol Laparosc.* 2002; 9:409-413
35. Angel E, Davis JR, Nagle RB. Immunohistochemical demonstration of placental hormones in the diagnosis of uterine versus ectopic pregnancy. *Am J Clin Pathol* 1985; 84:705-709.
36. Chokroverty M, Caballes RL, Gear PE. An unruptured tubal pregnancy at term. *Arch Pathol Lab Med.* 1986; 110:250-251
37. Deheragoda MG, Baithun S. Omental deposits of fetal parts as a sequel to salpingectomy for ruptured ectopic pregnancy. *Histopathology.* 2006; 49:426.
38. Burton JL, Lidbury EA, Gillespie AM, Tidy JA, Smith O, Lawry J *et al.* Over-diagnosis of hydatidiform mole in early tubal ectopic pregnancy. *Histopathology.* 2001; 38:409-417.
39. Blaustein A, Shenker L. Vascular lesions of the uterine tube in ectopic pregnancy. *Obstet Gynecol.* 1967; 30:551-555.
40. Milchgrub S, Sandstad J. Arias-Stella reaction in fallopian tube epithelium. A light and electron microscopic study with a review of the literature. *Am J Clin Pathol.* 1991; 95:892-895
41. Sheldon RS, Wilson RB, Dockerty MB. Serosal endometriosis of fallopian tubes. *Am J Obstet Gynecol* 1967; 99:882-884.
42. Rubin A, Czernobilsky B. Tubal ligation. A bacteriologic, histologic and clinical study. *Obstet Gynecol.* 1970; 36:199-203.
43. Kukreja P, Shetty KJ. Histopathological spectrum of lesions in fallopian tube. *IOSR-JDMS* 2017; 16(1):75-80.
44. Dahiya N, Sing S, Kalra R. *et al.* Histopathological changes associated with ectopic tubal pregnancy. *IJPSR* 2011; 2(4):929-933.
45. Shetty SK, Shetty AK. A clinical study of ectopic pregnancies in a tertiary care hospital of Mangalore. *Innov J Med Health Sci.* 2014; 4:305-309.
46. LI C, Meng CX, Zhao WH *et al.* Risk factors for ectopic pregnancy in women with planned pregnancy: A case-control study. *Eur J Obstet Gynecol Reprod Biol* 2014; 181:176-182.
47. Pratima Kujur, Shashikala Kosam, Anurag Gupta. Histopathological study of Spectrum of Lesions seen in surgically resected specimens of Fallopian Tube. *International Journal of Scientific Study* December 2016; 4:39-42.
48. Agarwal J, gupta JK. Female genital tuberculosis- a retrospective clinicopathologic study of 501 cases. *Indian J Pathol Microbiol.* 1993; 36(4):389-397.
49. Mehnaz Choudhary, Roopali Jandial. Histopathological Spectrum of lesions encountered in Fallopian Tube: A Study of 250 cases. *Indian Journal of Applied Research.* 2019; 9(11):1-3